

# got research?

405.773-6613/405 636-7951

## ► ICIO SMC study description May 2025

STUDY NAME	DZ SITE	STUDY Description	DESIGN		# PTS
NRG-GI008	Stage IIA Colon Cancer <b>Natera Kits provided by the Study</b>	<i>Colon Adjuvant Chemotherapy based on Evaluation of Residual Disease (Circulate-US)</i> <b>Cohort B – ctDNA positive</b> <b>Stratification – Intended Chemo (5 FU vs. Xeloda)</b> <b>Randomization</b> <b>Arm 3 – mFOLFOX 6 or CAPOX for 6 months</b> <b>A4m 4 – mFOLFIRINOX for 6 months</b>	<b>Cohort A</b> <b>ctDNA-neg</b> <b>Stratification- Stage IIIA vs IIIB</b> <b>Intended chemo (5FU vs. Xeloda)</b>	<b>After Randomization:</b> <b>Arm 1 FOLFOX6</b> for 3 – 6 months or CAPOX for 3 months. <b>Arm 2 –</b> Surveillance	
S2302	NSCLC	A Prospective Randomized Study of Ramucirumab + Pembrolizumab vs. standard of care for participants previously treated with Immunotherapy for Stage IV or recurrent NSCLC	<b>Arm A:</b> Investigator's SOC	<b>Arm B:</b> Ramucirumab + Pembrolizumab	
A081801	NSCLC	<i>NSCLC: ALCHEMIST-IO</i> <b>Must have EGFR, ALK, PD-LI IHC local testing with no identified arrangements. Must have completely resected dz IB (&gt;= 4cm), II or IIIA NSCLC with negative margins Patients with N2 Disease completely resected are eligible. However, patients with known N2 dz prior to surgery or ineligible.</b>	<b>Arm A:</b> 4 Cycles of Platinum doublet With Observation afterwards	<b>Arm B:</b> 4 Cycles of Platinum doublet with 17 cycles of Keytruda <b>Arm C:</b> 4 Cycles of Platinum doublet + Keytruda with 13 more cycles of Keytruda	1
EA5182	NSCLC	Randomized Phase III Study of Combination AZD9291 (Osimertinib) and Bevacizumab vs. AZD9291 (Osimertinib) Alone as First Line Treatment for Patients with Metastatic EGFR-Mutant Non Small Cell Lung Cancer (NSCLC)	<b>Arm A:</b> Osimertinib 80 mg PO Daily	<b>Arm B:</b> Osimertinib 80 mg Daily; Bevacizumab 15 mg kg IV every 3 weeks	
S2303	<b>Gastric and Esophageal Adenocarcinoma</b>	“Randomized Phase II/III Trial of 2nd Line Nivolumab + Paclitaxel + Ramucirumab versus Paclitaxel + Ramucirumab in Patients with PD-L1 CPS ≥ 1 Advanced Gastric and Esophageal Adenocarcinoma (PARAMUNE).”	<b>Arm 1:</b> Nivolumab, Paclitaxel, Ramucirumab	<b>Arm 2:</b> Paclitaxel, Ramucirumab	
CVAY73612301	Novartis	A phase III, randomized, double-blind study of Ianalumab (VAY736) vs. placebo in addition to first-line corticosteroids in primary immune thrombocytopenia	<b>Arm A:</b> Ianalumab 3mg/kg every 4 weeks (4 doses in total) + corticosteroids daily	<b>Arm B:</b> Ianalumab 9mg/kg every 4 weeks + corticosteroids daily <b>Arm C:</b> Placebo every 4 weeks	
ARGX-113-2402	ARGENX	A Phase 3, Multicenter, Randomized, Double-Blinded, Placebo-Controlled, Parallel-Arm Study Followed by an Open-Label Arm to Evaluate the Efficacy and Safety of Efgartigimod IV in Adult Participants With Primary Immune Thrombocytopenia	<b>Arm A:</b> Efgartigimod IV 10 mg/kg <b>Arm B:</b> Placebo	<b>After 24 weeks patients will receive Efgartigimod IV 10 mg/kg for 52 weeks</b>	